

Suicide Prevention Profile: Indicator List

Suicide data	Related risk factors	Related service contacts
County or UA	County or UA	County or UA
Suicide age-standardised rate 15+ years: per 100,000 (3 year average) (female)	Estimated prevalence of opiates and/or crack cocaine use: rate per 1,000 aged 15 - 64	Emergency Hospital Admissions for Intentional Self-Harm: Directly age-sex standardised rate per 100,000
Suicide age-standardised rate 15+ years: per 100,000 (3 year average) (persons)	Higher risk drinking: % of people drinking at increasing or higher risk levels	Adults in treatment at specialist drug misuse services: rate per 1000 population
Suicide age-standardised rate 15+ years: per 100,000 (3 year average) (male)	Long-term health problems or disability: % of people whose day-to-day activities are limited by their health or disability	Adults in treatment at specialist alcohol misuse services: rate per 1000 population
Persons years of life lost due to suicide age-standardised rate 15-74 years: rate per 10,000 persons (3 year average)	Self-reported well-being: % of people with a low satisfaction score	Successful completion of drug treatment - opiate users: % who do not re-present within 6 months
Female years of life lost due to suicide age-standardised rate 15-74 years: rate per 10,000 females (3 year average)	Self-reported well-being: % of people with a low worthwhile score	Successful completion of drug treatment - non-opiate users: % who do not re-present within 6 months
Male years of life lost due to suicide age-standardised rate 15-74 years: rate per 10,000 males (3 year average)	Self-reported well-being: % of people with a low happiness score	Successful completion of treatment for alcohol: % who do not re-present within 6 months
Female suicide crude rate 15-34 years: per 100,000 females (5 year average)	Self-reported well-being: % of people with a high anxiety score	Social care mental health clients receiving services during the year: Rate per 100,000 population
Female suicide crude rate 35-64 years: per 100,000 females (5 year average)	Prisoner population: Number	
Female suicide crude rate 65+ years: per 100,000 females (5 year average)	Children in the youth justice system: rate per 1,000 aged 10 - 18	CCG
Male suicide crude rate 15-34 years: per 100,000 males (5 year average)	Looked after children: Rate per 10,000 <18 population	Patients on QOF SMI register who have comprehensive care plan: % with plan
Male suicide crude rate 35-64 years: per 100,000 males (5 year average)	Children leaving care: Rate per 10,000 <18 population	Primary care assessment of depression: % of adults with a new diagnosis of depression who had a bio-psychosocial assessment
Male suicide crude rate 65+ years: per 100,000 males (5 year average)	Domestic abuse incidents: rate per 1,000 population	Access to IAPT services: People entering IAPT as % of those estimated to have anxiety/depression (in month)
	Relationship breakup: % of adults whose current marital status is separated or divorced	IAPT recovery: % of people who completed treatment who are "moving to recovery" (in month)
	Statutory homelessness: rate per 1000 households	People in contact with Adult Mental Health services: Rate per 100,000 population (end of quarter snapshot)
	People living alone: % of all households occupied by a single person	Mental Health Service users with crisis plans: % of people in contact with services with a crisis plan in place (end of quarter snapshot)
	Older people living alone: % of households occupied by a single person aged 65 or over	People on Care Programme Approach: Rate per 100,000 population (end of quarter snapshot)
	Unemployment: % of working age population	CPA adults in employment: % of people aged 18-69 on CPA in employment (end of quarter snapshot)
	long-term unemployment: % of working age population	CPA adults in settled accommodation: % of people aged 18-69 on CPA in settled accommodation
		Follow up after discharge: % of patients on CPA who were followed up within 7 days after discharge from psychiatric inpatient care (in quarter)
		Attendances at A&E for a psychiatric disorder: Rate per 100,000 population
	CCG	
	Number of people with SMI known to GPs: % on register	
	Number of adults with depression known to GPs: % on register	
	People estimated to have any common mental health disorder: Estimated % of population aged 16-74	