

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Updated May 2016



Key results for South West

Introduction

The Segment Tool has been developed by Public Health England's (PHE) Epidemiology and Surveillance team and provides information on the causes of death that are driving inequalities in life expectancy at local area level. Targeting the causes of death which contribute most to the life expectancy gap should have the biggest impact on reducing inequalities.

This document presents key results for South West and provides data tables and charts showing the breakdown of the life expectancy gap in 2012-14 for two comparisons:

1. The gap between the region as a whole and England as a whole.
2. The gap between the most deprived quintile of the region and the least deprived quintile of the region.

The tool, along with a user guide and technical document, can be downloaded from <http://fingertips.phe.org.uk/profile/segment>

Due to changes made in this version of the Segment Tool, the results presented here are not directly comparable with previous versions.

Interpreting the charts

Two types of chart are included below.

The scarf charts show, for each broad cause of death, the percentage contribution that it makes to the overall life expectancy gap between the areas selected. This could be between the region as a whole and England, or between the most deprived quintile of the region and the least deprived quintile of the region, depending on which comparison has been selected. If a cause shows a contribution of 0, this means that the cause of death does not make any contribution to the life expectancy gap.

The bar charts show, for a more detailed breakdown of causes of death, the years of life expectancy that would be gained or lost in the selected area, if it had the same mortality rates as the comparator area.

The tables show the percentage contributions and years of life gained for each cause of death. The number of deaths occurring in the area in 2012-14 are shown, and alongside, the number of excess deaths. Excess deaths are the number of 'extra' deaths that occur in the selected area because it has a higher mortality rate for that cause of death than the comparator area. If these deaths were prevented, then the contribution of that cause of death to the overall life expectancy gap would be eliminated. For some causes of death, there are no excess deaths in the selected area, and therefore no impact is made to the life expectancy gap. If this is the case the table shows a value of "..".

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Contextual information for South West

Information on inequalities in life expectancy between South West and England

	Male	Female
Life expectancy at birth in South West, 2012-2014	80.2	83.9
Life expectancy at birth in England, 2012-2014	79.6	83.2
Absolute gap in life expectancy between South West and England in years*	0.6	0.7

*A value below 0 indicates a lower life expectancy in the selected area compared to England.

Information on inequalities in life expectancy within South West

	Male	Female
Life expectancy at birth in the most deprived quintile of South West, 2012-2014	76.3	81.4
Life expectancy at birth in the least deprived quintile of South West, 2012-2014	83.0	85.8
Absolute gap in life expectancy between most deprived and least deprived areas within South West*	-6.7	-4.4

*A value below 0 indicates a lower life expectancy in the most deprived quintile compared with the least deprived quintile.

Sources:

- Life expectancies for regions - Office for National Statistics (ONS), life expectancy at birth for local areas in England and Wales
- Life Expectancies for deprivation quintiles within each local authority - Analysis by Public Health England Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Between area inequalities: Life expectancy gap between South West as a whole and England

Chart 1: Scarf chart showing the breakdown of the life expectancy gap between South West as a whole and England as a whole, by broad cause of death, 2012-2014

****This chart is not available as life expectancy is higher in South West than in England as a whole****




Footnote: Circulatory diseases includes coronary heart disease and stroke. Respiratory diseases includes flu, pneumonia and chronic obstructive airways disease. Digestive diseases includes alcohol-related conditions such as chronic liver disease and cirrhosis. External causes include deaths from injury, poisoning and suicide. Mental and behavioural includes dementia and Alzheimer's disease.

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Chart 2: Bar chart showing life expectancy years gained or lost if South West as a whole had the same mortality rates as England as a whole, by broad cause of death, 2012-2014



Footnote: Circulatory diseases includes coronary heart disease and stroke. Respiratory diseases includes flu, pneumonia and chronic obstructive airways disease. Digestive diseases includes alcohol-related conditions such as chronic liver disease and cirrhosis. External causes include deaths from injury, poisoning and suicide. Mental and behavioural includes dementia and Alzheimer's disease.

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates.

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Table 1: Breakdown of the life expectancy gap between South West as a whole and England as a whole, by broad cause of death, 2012-2014

Broad cause of death	Male				Female			
	Number of deaths in region	Number of excess deaths in region	Number of years of life gained/lost*	Contribution to the gap (%)	Number of deaths in region	Number of excess deaths in region	Number of years of life gained/lost*	Contribution to the gap (%)
Circulatory	21,877	-1,431	-0.23	++	23,033	-929	-0.16	++
Cancer	24,017	-1,245	-0.19	++	20,746	-1,436	-0.22	++
Respiratory	10,079	-1,319	-0.18	++	10,488	-1,969	-0.24	++
Digestive	3,496	-337	-0.07	++	3,866	-306	-0.07	++
External causes	3,566	94	0.04	++	2,424	74	0.04	++
Mental and behavioural	5,532	-376	-0.03	++	11,246	-643	-0.07	++
Other	8,225	95	-0.01	++	11,556	483	0.01	++
Deaths under 28 days	274	-1	0.00	++	171	-45	-0.04	++
Total	77,065			++	83,530			++

Counts of deaths have been adjusted take account of changes in ICD-10 coding made in 2014. This means totals may differ between tables. Further details can be found in the technical document.

* A positive figure indicates that life expectancy years would be gained if the region had the same mortality rate as England as a whole (ie the mortality rate in the region for the cause is higher than England as a whole). A negative figure indicates that life expectancy years would be lost if the region had the same mortality rate as England as a whole.

.. The calculated contribution to the gap is negative


++ The figures have not been calculated as life expectancy is higher than England

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates.

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Chart 3: Bar chart showing life expectancy years gained or lost if South West as a whole had the same mortality rates as England as a whole, by detailed cause of death, 2012-2014



Footnote: See table below for details of cause groupings. Details of causes of death included in the 'other' category can be found in the Technical Document. Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates.

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Table 2: Table showing life expectancy years gained or lost if South West as a whole had the same mortality rates as England as a whole, by detailed cause of death, 2012-2014

Broad cause of death	Detailed cause of death	Male			Female		
		Number of deaths in region	Number of excess deaths in region	Number of years of life gained/lost*	Number of deaths in region	Number of excess deaths in region	Number of years of life gained/lost*
Circulatory	Coronary Heart Disease	11,326	-1,031	-0.17	8,117	-617	-0.10
	Stroke	4,825	-37	-0.01	7,845	565	0.05
	Other circulatory	5,732	-356	-0.05	7,115	-871	-0.10
Cancer	Lung cancer	4,761	-848	-0.12	3,538	-1,001	-0.14
	Other cancers	19,256	-396	-0.07	17,208	-435	-0.08
Respiratory	Pneumonia	3,989	-145	-0.03	5,090	-286	-0.04
	Chronic obstructive airways disease	4,058	-629	-0.08	3,400	-977	-0.12
	Other respiratory disease	2,043	-472	-0.07	2,016	-616	-0.07
Digestive	Chronic liver disease including cirrhosis	1,012	-179	-0.04	539	-138	-0.04
	Other digestive	2,459	-153	-0.03	3,327	-169	-0.03
External	Suicide	1,182	91	0.02	433	102	0.04
	Other external	2,384	2	0.02	1,995	-34	-0.01
Mental and behavioural	Dementia & Alzheimer's disease	5,358	-391	-0.04	11,122	-646	-0.07
	Other mental and behavioural disorders	175	15	0.01	122	3	0.00
Other	Infectious and parasitic diseases	754	-35	-0.01	884	-67	-0.02
	Urinary conditions	1,068	-144	-0.02	1,491	-245	-0.03
	Ill defined conditions	1,248	244	0.04	3,670	752	0.08
	Diabetes	820	-33	-0.01	846	-68	-0.01
	Other	4,332	72	-0.01	4,626	130	0.00
< 28 days	Deaths under 28 days	274	-1	0.00	171	-45	-0.04
Total		77,055			83,554		

Counts of deaths have been adjusted take account of changes in ICD-10 coding made in 2014. This means totals may differ between tables. Further details can be found in the technical document.

* A positive figure indicates that life expectancy years would be gained if the region had the same mortality rate as England as a whole (ie the mortality rate in the region for the cause is higher than England as a whole). A negative figure indicates that life expectancy years would be lost if the region had the same mortality rate as England as a whole.


Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates.

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Within area inequalities: Life expectancy gap between the most deprived quintile and least deprived quintile of South West

Chart 4: Scarf chart showing the breakdown of the life expectancy gap between South West most deprived quintile and South West least deprived quintile, by broad cause of death, 2012-2014




Footnote: Circulatory diseases includes coronary heart disease and stroke. Respiratory diseases includes flu, pneumonia and chronic obstructive airways disease. Digestive diseases includes alcohol-related conditions such as chronic liver disease and cirrhosis. External causes include deaths from injury, poisoning and suicide. Mental and behavioural includes dementia and Alzheimer’s disease.

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Chart 5: Bar chart showing life expectancy years gained or lost if South West most deprived quintile had the same mortality rates as South West least deprived quintile, by broad cause of death, 2012-2014



Footnote: Circulatory diseases includes coronary heart disease and stroke. Respiratory diseases includes flu, pneumonia and chronic obstructive airways disease. Digestive diseases includes alcohol-related conditions such as chronic liver disease and cirrhosis. External causes include deaths from injury, poisoning and suicide. Mental and behavioural includes dementia and Alzheimer’s disease.

Analysis by Public Health England’s (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Table 3: Breakdown of the life expectancy gap between South West most deprived quintile and South West least deprived quintile, by broad cause of death, 2012-2014

Broad cause of death	Male				Female			
	Number of deaths in most deprived quintile	Number of excess deaths in most deprived quintile	Number of years of life gained/lost*	Contribution to the gap (%)	Number of deaths in most deprived quintile	Number of excess deaths in most deprived quintile	Number of years of life gained/lost*	Contribution to the gap (%)
Circulatory	4,648	1,945	1.64	27.8	4,655	1,288	1.06	26.4
Cancer	4,663	1,364	1.21	20.4	4,039	850	0.85	21.2
Respiratory	2,243	1,049	0.80	13.6	2,411	990	0.80	19.9
Digestive	937	520	0.56	9.4	913	355	0.36	9.0
External causes	984	540	0.83	14.0	553	226	0.33	8.1
Mental and behavioural	1,017	230	0.17	2.8	2,054	300	0.17	4.3
Other	1,624	520	0.60	10.1	2,161	265	0.40	10.0
Deaths under 28 days	81	34	0.11	1.8	49	13	0.05	1.2
Total	16,198			100	16,834			100

Counts of deaths have been adjusted take account of changes in ICD-10 coding made in 2014. This means totals may differ between tables. Further details can be found in the technical document.

* A positive figure indicates that life expectancy years would be gained if the most deprived quintile of the region had the same mortality rate as the least deprived quintile (ie the mortality rate in the most deprived quintile for the cause is higher than that of the least deprived quintile). A negative figure indicates that life expectancy years would be lost if the most deprived quintile in the region had the same mortality rate as the least deprived quintile.

.. The calculated contribution to the gap is negative


The figures have not been calculated as life expectancy in the most deprived quintile is higher than life expectancy in the least deprived quintile

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Chart 6: Bar chart showing life expectancy years gained or lost if South West most deprived quintile had the same mortality rates as South West least deprived quintile, by detailed cause of death, 2012-2014



Footnote: See table below for details of cause groupings. Details of causes of death included in the 'other' category can be found in the Technical Document. Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015

THE SEGMENT TOOL

SEGMENTING LIFE EXPECTANCY GAPS BY CAUSE OF DEATH

Table 4: Table showing life expectancy years gained or lost if South West most deprived quintile had the same mortality rates as South West least deprived quintile, by detailed cause of death, 2012-2014

Broad cause of death	Detailed cause of death	Male			Female		
		Number of deaths in most deprived quintile	Number of excess deaths in most deprived quintile	Number of years of life gained/lost*	Number of deaths in most deprived quintile	Number of excess deaths in most deprived quintile	Number of years of life gained/lost*
Circulatory	Coronary Heart Disease	2,535	1,175	0.99	1,685	534	0.42
	Stroke	906	274	0.21	1,519	313	0.24
	Other circulatory	1,206	494	0.40	1,461	445	0.38
Cancer	Lung cancer	1,168	611	0.50	908	483	0.46
	Other cancers	3,495	754	0.68	3,131	368	0.38
Respiratory	Pneumonia	777	249	0.19	1,021	262	0.18
	Chronic obstructive airways disease	1,070	677	0.49	1,004	670	0.53
	Other respiratory disease	401	130	0.11	398	74	0.08
Digestive	Chronic liver disease including cirrhosis	358	249	0.31	145	88	0.11
	Other digestive	571	265	0.24	768	267	0.24
External	Suicide	327	174	0.28	111	51	0.10
	Other external	657	365	0.54	443	177	0.23
Mental and behavioural	Dementia & Alzheimer's disease	947	177	0.10	2,014	279	0.14
	Other mental and behavioural disorders	70	52	0.07	39	21	0.03
Other	Infectious and parasitic diseases	186	97	0.09	207	80	0.08
	Urinary conditions	227	113	0.07	319	115	0.08
	Ill defined conditions	265	84	0.16	593	-100	0.00
	Diabetes	168	67	0.06	192	67	0.06
	Other	778	158	0.21	840	98	0.19
< 28 days	Deaths under 28 days	81	34	0.11	49	13	0.05
Total		16,194			16,847		

Counts of deaths have been adjusted take account of changes in ICD-10 coding made in 2014. This means totals may differ between tables. Further details can be found in the technical document.

* A positive figure indicates that life expectancy years would be gained if the most deprived quintile of the region had the same mortality rate as the least deprived quintile (ie the mortality rate in the most deprived quintile for the cause is higher than that of the least deprived quintile). A negative figure indicates that life expectancy years would be lost if the most deprived quintile in the region had the same mortality rate as the least deprived quintile.

Analysis by Public Health England's (PHE) Epidemiology and Surveillance team based on ONS death registration data, and mid year population estimates, and DCLG Index of Multiple Deprivation, 2015